



idena

THE ANIMAL NUTRITION, NATURALLY INNOVATIVE

# FORCIX SW

THE NATURAL ALTERNATIVE

ECHV  
Technology



## FOR THE ANIMAL

Intestinal integrity of piglets



## FOR THE FARMER

Easy to use



## FOR THE CONSUMER

Healthier food

[idena.fr/en](http://idena.fr/en)

Distributed in Australia by BEC Feed Solutions  
Phone: 1300 884 593 - [becfeedsolutions.com.au](http://becfeedsolutions.com.au)



# FORCIX SW

At birth, piglets have a sterile gut. **Soon after, a diverse microbial community originating from the sow and its environment populates the gut.**

Parasites such as coccidia have been shown to affect the integrity of the piglets' intestinal mucosa, leading to incidence of dysbiosis at day 10 to 15 of age. The affected piglets have poor growth rate due to malabsorption and increased susceptibility to other disease challenges.


**FORCIX SW is a combination of several natural essential oils and plant extracts selected with the phytogramme® process and field-tested.**

Distributed to the sow at pre-farrowing, **FORCIX SW** helps in management of the impact of protozoa (Eimeriidae family, type Eimeria and Cytoisospora) and promotes growth in young piglets.



## FORCIX SW effect on diarrhoea of piglet at 8 days of age


Source : Comparative trial, Australia, 2020



\* positive control = Toltrazuril 1ml per piglet at 3 days of age

## FORCIX SW effect on piglet growth during lactation

Source : Comparative trial, Australia, 2020



\* positive control = Toltrazuril 1ml per piglet at 3 days of age

|                                  | Toltrazuril (T1) | FORCIX SW (T2) | P-Value |
|----------------------------------|------------------|----------------|---------|
| Average ADG during lactation (g) | 237              | 253            | < 0.005 |

Significantly higher ADG in **FORCIX SW** group (+16g/day)

**In this trial, the usage of FORCIX SW led to improvement of weaning weight by 336g/piglet.**

## INSTRUCTION OF USE

**FORCIX SW is applied as a single application of 20ml in the sow feed , 5 to 6 days before farrowing.** For gilts sourced externally, 20 ml of **FORCIX SW** should be given during quarantine period.

Keep away from moisture and light, in a fresh, ventilated place.



THE ANIMAL NUTRITION, NATURALLY INNOVATIVE